CHEM 244 Organic Chemistry II Points ______ (10 max)

Worksheet #2a: January 29, 2025. Complete the following worksheet by collaborating with the students in your group. You can use a text book or your lecture video notes. You must work together, with the names of all students included on **ONE** sheet and turned in for a group grade.

PLEASE NOTE – Videos 12-3 and 12-4 require TWO class days of worksheets (WS-2a and WS-2b). Our class on Friday we will have worksheet 2b (no new videos), and we will start project work next week.

(1) Grignard Reaction True/False. Are the following statements about Grignard Reaction True or False.

- The Grignard Reaction is a 3-step sequence that requires an alkyl halide, Mg, ether solvent, an "oxygen containing compound" (aldehyde, ketone, ester or epoxide), and aqueous acid.
- _____ The Grignard Reaction has a carbocation intermediate.

_____ The Grignard <u>reagent</u> is best characterized as (1) a strong base, and (2) a good nucleophile.


_____ The final product can be a 1°, 2°, or 3° alcohol.

_____ Water is the typical solvent when making the Grignard reagent.

(2) **Reactions used in the Grignard synthesis.** Fill in the boxes to give the correct structure of the alcohol product in the <u>paired reactions</u> shown below. In the first reaction in each pair, the "dots" represent where the new C-C bond will form and are to guide you in drawing the correct alcohol product. In the second reaction in each pair, the dots are not included.


I. Grignard Pair #1: Reaction of a Grignard Reagent with an Aldehyde.

(a) When a Grignard Reagent reacts with an <u>aldehyde</u>, will the product alcohol be 1°, 2°, or 3°? Answer: _____




II. Grignard Pair #2: Reaction of an Alkyl Halide with a Ketone.


(d) When a Grignard Reagent reacts with a ketone, will the product alcohol be 1°, 2°, or 3°? Answer: _____




III. Grignard Pair #3: Reaction of a Grignard Reagent with an Ester. NOTE: TWO molecules of the Grignard Reagent must add to ONE molecule of an ester.

(g) When a Grignard Reagent reacts with an <u>ester</u>, will the product alcohol be 1°, 2°, or 3°? **Answer:**_____





(j) When a Grignard Reagent reacts with an epoxide, will the product alcohol be 1°, 2°, or 3°? Answer: _



(3) **REVIEW:** Alcohol Nomenclature.

- <u>If a name is given</u> draw an accurate zig-zag structure (remember to use wedge and dash bonds for all chiral carbons).
- If a structure is drawn, give an accurate IUPAC name (don't forget to assign configurations using the R/S prefix).

(2S, 4S)-1-cyclopropyl-4-methyl-2-hexanol