E.F. Devenay / BSC Physics, PH458 E&M II Chpt 8 Griffiths

Note Title 4/6/2005

$\vec{E}, \vec{B}$ & $\vec{\mathcal{E}}$ in FIELDS

Wow! we've got Maxwell's Equations

$$\vec{\nabla}\cdot\vec{E} = \frac{\rho_f}{\varepsilon_0} + 0$$

$$\vec{\nabla}\times\vec{E} = 0 + -\frac{d\vec{B}}{dt}$$

$$\vec{\nabla}\cdot\vec{B} = 0 + 0$$

$$\vec{\nabla}\times\vec{B} = \mu_0\vec{J}_f + \mu_0\varepsilon_0\frac{d\vec{E}}{dt}$$

static source dynamic sources

of $\vec{\nabla}\cdot$ & $\vec{\nabla}\times$ $\vec{E}$ & $\vec{B}$

<u>Completely</u> describe Vector Fields!

(recall $\vec{\nabla}\cdot \propto$  $\hat{r}$

& $\vec{\nabla}\times \propto$  $\propto \hat{\theta}, \hat{\phi}$

so $\vec{\nabla}$ & $\vec{\nabla}$ span 3-D of those vector fields

Now we have the ability to look @ problems from 2 equiv reps



work done in building current

$= \frac{1}{2} L I^2$

Energy stored in field

$= \frac{1}{\mu_0} \int |B|^2 d\tau$

use what ever is best suited for problem!

Big Idea: If I see a problem w/ Electricity & magnetism, if it is useful, you can cast the problem entirely in terms of fields using Maxwell's Equations!

Also: if you don't think about "fields", you run into problems!

(ie Conservation laws Need "fields" to Conserve!)

So, chapt 8 Griff = properties of Fields!

Lets Begin by considering some conservation inconsistencies!

1) pg 307 Griff ex: 7.8



$\mathcal{E}\vec{\mathcal{P}}_{net} = \frac{d\vec{U}}{dt}$

$0 = \frac{d\vec{U}}{dt}$

So $\vec{L}$ = conserved

But here, $L_i = 0$

$L_f \neq 0$

?

2.) Why you Toaster Sparks?



w/ $I=0$,
where did Energy come from?

3.)



$F_{mag\,b} = q_b(\vec{V}_b \times \vec{B}_a) = (-\hat{x})$

$F_{mag\,a} = q_a(\vec{V}_a \times \vec{B}_b) = (+\hat{y})$

Note: $\vec{E}$ & $\vec{B}$ not static, but indeed directions OK Guess

so

$|F_{m_b}| = |F_{m_a}|$

but not opposite direction!

In violation of Newton's 3rd law

Now might not seem too big a deal BUT conservation of linear moment

Very much depends on cancelation of internal forces



$$\sum \vec{F} = m\ddot{x} = \frac{d}{dt}(\vec{p})$$

$$\underbrace{0}_{\text{is all internal}} = \frac{d\vec{p}}{dt} \quad \text{or} \quad \vec{p} = \text{constant!}$$

"conserved."

How do reconcile these "violations" to

Conservation of angular momentum, $\vec{L}$

" " energy, E

" " linear momentum, $\vec{p}$

The Answer of course is .....

The Field "Carries" $\vec{L}$, E & $\vec{p}$ to conserve!

Here is Basic idea again.....

Continuity Equation (general)

$$\vec{\nabla}\cdot\vec{J} = -\underbrace{\frac{\partial(\text{conserved stuff})}{\partial t}}$$

if you have stuff that is conserved & it "disappears" (ie $\frac{\partial}{\partial t} \neq 0$)

$\Rightarrow$'s a Field is "Sourced"

so creates a Field Current & this carries the conserved quantities!

ex: Color charge = conserved = strong nuclear Force so if quark color changes, A Field current must be created that conserves color



$|i\rangle$ = red

$|f\rangle = g + \bar{g} + r = r$

gluon color field

So in order to conserve

$\vec{L}$, E & $\vec{p}$ you need fields

or

$\Rightarrow$ Since $\vec{L}$, E & $\vec{p}$ MUST be

conserved based on symmetry

arguments (ie no perfect

symm	cons
$x_0 \Rightarrow$	$\vec{p}$
$\theta_0 \Rightarrow$	$\vec{L}$
$t_0 \Rightarrow$	E

you

MUST

HAVE

FIELDS $\therefore$ HAVE FORCES!

ie standard Model

OK! $\vec{E}$ & $\vec{B}$ fields "carry" $\vec{L}$, $\vec{p}$ & E

and combine to conserve $\vec{L}$, $\vec{p}$ & E

Final results on $\vec{E}$ & $\vec{B}$ fields from Chpt 8

Energy E & B $= u_{EM} = \frac{1}{2}\left(\varepsilon_0 E^2 + \frac{1}{\mu_0} B^2\right) = \frac{\text{energy}}{\text{volume}}$

momentum: $\vec{p}_{E\&M} = \varepsilon_0 (\vec{E} \times \vec{B}) = \frac{\text{momentum}}{\text{volume}}$

angular momentum $= \vec{r} \times \vec{p}_{E\&M} = \varepsilon_0 \left[\vec{r} \times (\vec{E} \times \vec{B})\right]$

Carried by FIELDS

Great results & will need ∞-ly. Examples

$\Longrightarrow$

Why need $\vec{P}$, $\vec{B}$ & E of fields!

ex: atom interacting with light



$$\hat{H}_0 = E_K + E_P = \frac{\hat{P}^2}{2m} - \frac{Ze^2}{r}$$

Solve $i\hbar \frac{\partial \Psi(t,\vec{r})}{\partial t} = \hat{H}_0 \Psi(t,\vec{r})$

& get $\Psi_{100} = e^{-i\frac{E_n}{\hbar}t} R_n(r)\, Y_\ell^m(\theta,\phi)$

now these = stationary!

They don't change in time

See of Course Townsend Chapt 14 GREAT



stays Forever!

Now is couple to ⟹



self prop $\vec{E}$ & $\vec{B}$ field

$\vec{\nabla} \times \vec{E} = -\frac{d\vec{B}}{dt}$

$\vec{\nabla} \times \vec{B} = \mu_0 \epsilon_0 \frac{d\vec{E}}{dt}$

$= \gamma$

Transverse real photon

need $\hat{H} = \hat{H}_{0\,atom} + \hat{H}_{E\&M}$ $\sim \int \left[\epsilon_0 |E|^2 + \frac{1}{\mu_0} \vec{B}^2\right] d\tau$

Now atom can decay cause it is coupled to γ field

So



Now

$$E_i = E_s$$
$$\vec{p}_i = \vec{p}_s$$
$$\vec{L}_i = \vec{L}_s$$

atom = atom + field

So $\vec{E}$ & $\vec{M}$ fields

$$u_{E\&M} = \frac{1}{2}\left(\varepsilon_0 E^2 + \frac{1}{\mu_0} B^2\right)$$

$$\vec{p}_{E\&M} = \varepsilon_0 (\vec{E} \times \vec{B}) \qquad = \mu_0 \varepsilon_0 \vec{S}$$

$$\vec{l}_{E\&M} = \varepsilon_0 [\vec{r} \times (\vec{E} \times \vec{B})] \qquad = \mu_0 \varepsilon_0 [\vec{r} \times \vec{S}]$$

Introduced $\vec{S}$ & $\vec{S}$ = Poynting Vector & plays Big role.....

Poynting Vector:

Start w/ static charge distribution & $\vec{E}$'s & $\vec{B}$'s



Now let charges move abit .....

? = how much work is done by fields on these charges

well $dw = \vec{F} \cdot d\vec{l}$

$$dw = \vec{F}\cdot d\vec{l} = q(\vec{E} + \vec{v}\times\vec{B})\cdot\vec{v}\,dt$$

$$dw = q\vec{E}\cdot\vec{v}\,dt$$

* again $\vec{B}$'s do not work!

since

$$dq = \rho\, d\tau$$

& $\rho\vec{v} = \vec{J}$

Then

$$\frac{dw}{dt} = \vec{E}\cdot q\vec{v}\,dt$$

$$= \vec{E}\cdot \rho\, d\tau\, \vec{v}$$

$$\frac{dw}{dt} = \vec{E}\cdot\vec{J}\,d\tau$$

or?

$$\frac{dw}{dt} = \int_{\mathcal{V}} (\vec{E}\cdot\vec{J})\,d\tau$$

This is cool, but $\vec{J}$ .... Huh, represent in terms of pure fields .....

use Maxwell's equat's

Lots of steps.... Some magic

pg 347

get ....

$$\frac{dW}{dt} = -\frac{d}{dt}\int_{V} \frac{1}{2}\left(\varepsilon_0 E^2 + \frac{1}{\mu_0}B^2\right) d\tau - \frac{1}{\mu_0}\oint_{surf} (\vec{E}\times\vec{B})\cdot d\vec{a}$$

So .... $\frac{dW}{dt}$ = power

So RHS = power supplied by FIELDS!

we know $\frac{1}{2}\left(\varepsilon_0 E^2 + \frac{1}{\mu_0}B^2\right) = \frac{\text{Energy}}{\text{Volume}} = u_{E\&M}$

Introduce $\vec{S} = \frac{1}{\mu_0}(\vec{E}\times\vec{B})$ must $= \frac{\text{Energy}}{(\text{time})(\text{Area})} = \frac{\text{Power}}{\text{Area}}$

= poynting Vector!

So

$$\boxed{\frac{dW}{dt} = -\frac{du_{E\&M}}{dt} - \oint_{surf} \vec{S}\cdot d\vec{a}}$$

Now what does this

$$\boxed{\frac{dw}{dt} = \frac{-dU_{EM}}{dt} - \oint \vec{S} \cdot d\vec{a}}$$

mean?

Here is the deal



= Changes of energy density of field to make things like current & voltage = power

Stay w/in volume

$\mathcal{E}_{ms} = -\frac{d\Phi_B}{dt}$

+

Power that leaves the surface ie $\vec{E}$ & $\vec{M}$ waves = photons!

So, in chpt 9 on $\vec{E}$ & $\vec{M}$ waves,

Starting pts are

1.) $u_{E\&M} = \frac{1}{2}\left(\varepsilon_0 E^2 + \frac{1}{\mu} B^2\right) = \frac{\text{energy}}{\text{vol}}$

&

2.) $\vec{S} = \frac{1}{\mu_0}\left(\vec{E} \times \vec{B}\right) = \frac{\text{energy}}{(\text{Area})(\text{time})} = \frac{\text{power}}{\text{Area}}$

H.W.:

GRIFFS

Example problems:

Ex: 7.8



where did $\vec{L}$ come from

Ex: 8.1 using $\vec{S}$

Ex: 8.3 $\vec{p}_{E\&M}$ → hidden momentum

Ex: 8.4 $\vec{l}_{E\&M}$ → no hidden momentum