

time-indep - Schro

2 Band System \Rightarrow 1) ∞ Square well
2) Harmonic Oscillator

1) ∞ Square well.

We've used this case already as
launching pad for really the entire
"computing" aspects of the QM formalism,
--- i.e. "shut up & calculate"
to get observables
 $\langle X \rangle \pm \Delta X$ for example

to do that, we just used the soln
to the ∞ well

$$\hat{H}\Psi(x,t) = i\hbar \frac{\partial \Psi(x,t)}{\partial t}$$

$$\Psi(x,t) = \begin{cases} \sqrt{\frac{2}{a}} \sin\left(\frac{\pi x}{a}\right) e^{-\frac{i\hbar\pi^2 t}{2ma^2}} & 0 \leq x \leq a \\ 0 & \text{otherwise.} \end{cases}$$


Now we will get this solus explicitly

by solving for $\Psi(x,t) = \Psi(x) e^{-i\frac{E_F}{\hbar}t}$

↑
from $\hat{H}\Psi(x) = E_F\Psi(x)$

So again we will find energy eigenstates!

Recall of square well = ideal case... but we will use it for \approx Quantitative results and for Qualitative feel of real problems.





1) real, true energies might be shifted slightly

2) ψ does leak a bit in real case

Also can expect that \approx to real is not as good when $E \approx V_0$

ψ_n as n gets Big!

*** 1-D**
 So ∞ square well = Bound i.e. \square by (∞)
 potential on 2 sides



Biological detectors need source of ψ_e = Quantum "dot" transistors, lasers, very hot tuning these devices


IS Sandwich in all 3 directions

$\psi_e = \psi_e$ = Quantum "dot"

$\psi_e = \psi_e$ = Quantum "dot"

$\psi_e = \psi_e$ = Quantum "dot"

$\hat{H}\psi = E_T\psi \Rightarrow \psi'' + A\psi = 0 ; A = \frac{2m}{\hbar^2}(E_T - V)$



$\psi(x) = C_1 e^{\sqrt{A}x} + C_2 e^{-\sqrt{A}x}$

$\psi_{II} \& \psi_{III} = \text{graph of } e^{\sqrt{A}x} + e^{-\sqrt{A}x}$

Blows up, can't normalize ψ so

$\psi(x) = C_2 e^{-\sqrt{A}x}$

$\sqrt{A} = \sqrt{\frac{2m}{\hbar^2}(E_T - \infty)} = \infty$

$\psi_{II} \& \psi_{III} = C_2 e^{-\infty} = 0$

No Penetration into ∞ potential well!

I guess we should reserve tunneling for "thru"

tunneling $\rightarrow 0$ & a potential \rightarrow this should be no penetration

Region I $E > V$ ($V=0$) : $\therefore \sqrt{A} = \frac{\sqrt{2mE}}{\hbar}$


So

$$\psi_I(x) = C_1 e^{i\sqrt{A}x} + C_2 e^{-i\sqrt{A}x}$$

OR

$$\psi_I = C_1 \cos(\sqrt{A}x) + C_2 \sin(\sqrt{A}x)$$

\Rightarrow easier to use this equivalent soln!



$$\psi_I(x) = C_1 \cos\left(\frac{\sqrt{2mE}}{\hbar} x\right) + C_2 \sin\left(\frac{\sqrt{2mE}}{\hbar} x\right)$$

Now all we have to $\&$ make sure that ψ meets the BC's.

* Note ψ $\&$ $\frac{d\psi}{dx}$ do not have to be smooth from $\pm \epsilon$ about $x=0$ $\&$ $x=a$. Outside of these regions, $\psi=0 \Rightarrow$ period. So all that we need to do is make sure $\psi(0) = \psi(a) = 0$

OK, lets do it

$$\text{ii) } \left. \begin{aligned} \psi(0) &= C_1 \cos(0) + C_2 \sin(0) = 0 \\ C_1 &= 0 \end{aligned} \right\} \text{ Done: } C_1 = 0 \text{ already}$$

$\psi(x) = C_2 \sin\left(\frac{\sqrt{2mE}}{\hbar} x\right)$ so far. Next

$$\psi(x=a) = C_2 \sin\left(\frac{\sqrt{2mE}}{\hbar} x\right) = 0$$

well $C_2 = 0$ is possibility but then

$\psi(x) = 0$ doesn't help so

So $C_2 \neq 0$
must be

$$\sin\left(\frac{\sqrt{2mE}}{\hbar} x\right) = 0$$

This is true for, i.e.,

$$\frac{\sqrt{2mE}}{\hbar} x = 0, \pi, 2\pi, 3\pi, \dots = n\pi$$

$$\frac{\sqrt{2mE}}{\hbar} = n\pi, \quad (n=0, 1, 2, 3, \dots)$$

$= 0$ or $E = 0$

so $\psi(x) = C_2 \sin(0) = 0$

doesn't help
either to have

$\psi(x)$ that vanishes
everywhere!

so $\frac{\sqrt{2mE}}{\hbar} = n\pi$ for $n=1, 2, 3, \dots$

or $E_T = \frac{h^2 \pi^2}{2ma^2} n^2$; $n=1, 2, 3, \dots$

↓
call

$E_n \Rightarrow$'s E_{TOT} for state $n = \frac{h^2 \pi^2}{2ma^2} n^2$

$n=1, 2, 3, \dots$

lets think:

$n=0$, no wave function, $\psi_0^* \psi = 0 =$ no particle

$n=1$

$E_1 = \frac{h^2 \pi^2}{2ma^2}$

$=$ lowest possible energy of the particle \equiv zero

↓
↓
↓

next possible ψ_n is ψ_2 w/

$E_2 = E_1 (2^2)$

! Then

$E_3 = E_1 (3^2)$

↓
 $E_n = E_1 (n^2)$

or energy of system:

IT is there, can't be any less than this!

QUANTUM Says: Particle energies are Quantized exist only in discrete states w/ energies that jump,

In between, for energies other than quantized in these steps, The particle $\psi=0$ & or doesn't "Exist"

Cool.... but back to $\psi(x) = ?$

well so far

$$\psi_{II}(x) = C_2 \sin\left(\frac{\sqrt{2mE}}{\hbar} x\right)$$

but $E_T = E_n$ so must have distinct ψ_n for each ψ_n

$$\psi_n(x) = C_2 \sin\left(\frac{\sqrt{2mE_n}}{\hbar} x\right)$$

$$= C_2 \sin\left(\frac{\sqrt{2m \left(\frac{n^2 \pi^2 \hbar^2}{4ma^2}\right)}}{\hbar} x\right)$$

$$\psi_n(x) = C_2 \sin\left(\frac{n\pi x}{a}\right)$$

What's left? Born's ... probabilistic requirement

$$\int_{-\infty}^{+\infty} \psi_n^*(x) \psi_n(x) dx = 1 \quad ; \quad \text{Let } C_2 = \text{norm constant}$$

$$|C_2|^2 \int_0^a \sin^2 \frac{n\pi x}{a} dx = 1$$

note limits

we've done a million times now! * need

$$\cos(\alpha \pm \beta) = \cos\alpha \cos\beta \mp \sin\alpha \sin\beta$$

$$\text{Then } \sin^2 \alpha + \cos^2 \alpha = 1$$

or

$$|C_2|^2 \frac{a}{2} = 1$$

$$C_2 = \pm \sqrt{\frac{2}{a}}$$

we know Q.M.


$(-1) = e^{i\pi} = \text{arbitrary phase, no effect on expectation}$

So choose $c_2 = +\sqrt{\frac{2}{a}}$

$$\Psi_n(x) = \begin{cases} \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right) & ; 0 \leq x \leq a \\ 0 & \text{otherwise} \end{cases}$$

where $E_{n \text{ or } n} = \frac{\hbar^2 \pi^2}{2m a^2} n^2 ; n=1, 2, 3 \dots \infty$

lets look @ $\Psi_n(x)$ & $\Psi_n^* \Psi_n$




* recall idea of $n = \# \text{ of } \frac{1}{2} \lambda = \text{standing wave condition.}$

This gives most in So: $E_n, \Psi_n, \Psi_n^* \Psi_n$

For $n \rightarrow \text{Big}$, $E_n = E_1 (n^2) = \text{BIG}$


$$\Psi_n \text{ & } \Psi_n^* \Psi_n$$



and as a particle...

① Tennis ball $\rightarrow \Psi_n$ as $n \rightarrow \infty$


$\&$ see that $\Psi_n^* \Psi_n = \text{Constant}$



equally likely
 \rightarrow
Sound classical
Ball anywhere!

* Note: Ball does not speed up or slow down on collisions w/ wall \therefore does not spend more time there.

* Note: moving to harmonic oscillator potential



$$V(x) = \frac{1}{2} kx^2$$

$\&$ clearly here
ball does slow
down @ $\pm a$

So should have
higher prob of
being found here!